



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Vanadium – a re-emerging environmental hazard

Citation for published version:

Watt, J, Burke, IT, Edwards, RA, Malcolm, HM, Mayes, WM, Olszewska, J, Pan, G, Graham, M, Heal, K, Rose, NL, Turner, SD & Spears, BM 2018, 'Vanadium – a re-emerging environmental hazard', *Environmental Science and Technology*, vol. 52, pp. 11973-11974. <https://doi.org/10.1021/acs.est.8b05560>

Digital Object Identifier (DOI):

[10.1021/acs.est.8b05560](https://doi.org/10.1021/acs.est.8b05560)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Environmental Science and Technology

Publisher Rights Statement:

© 2018 American Chemical Society

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Vanadium – a re-emerging environmental hazard**

2
3 James A.J. Watt^{1,6}, Ian T. Burke², Ron A. Edwards³, Heath M. Malcolm¹, William M. Mayes⁴
4 Justyna P. Olszewska¹, Gang Pan⁵, Margaret C. Graham⁶, Kate V. Heal⁶, Neil L. Rose⁷, Simon
5 D. Turner⁷, and Bryan M. Spears¹

6 ¹ Centre for Ecology & Hydrology, Penicuik, Midlothian, EH26 0QB, UK (*contact
7 spear@ceh.ac.uk)

8 ² Earth Surface Science Institute, School of Earth and Environment, University of Leeds,
9 Leeds, LS2 9JT, UK

10 ³ Craigencalt Rural Community Trust, Kinghorn, Fife, KY3 9YG, UK

11 ⁴ School of Environmental Sciences, University of Hull, Hull, HU6 7RX, UK

12 ⁵ School of Animal, Rural and Environmental Sciences, Nottingham Trent University,
13 Nottinghamshire, NG25 0QF, UK

14 ⁶ School of GeoSciences, University of Edinburgh, Edinburgh, EH9 3FF

15 ⁷ Environmental Change Research Centre, Department of Geography, University College
16 London, London, WC1E 6BT, UK

Introduction

Vanadium (V) is a contaminant which has been long confined to the annals of regulatory history. This follows the reduction of its historical primary source (fossil fuel emissions) since the 1970s (e.g. by 80% in the UK). However, V is quickly becoming an important strategic resource which promises its return to environmental prominence because of changing industrial practices and emerging waste streams. We discuss below: (i) what makes V a re-emerging environmental and human health hazard of global interest, (ii) the knowledge gaps that currently restrict prediction of environmental effect and mitigation, and (iii) opportunities for the community to address these gaps towards reducing the risk of an impending environmental hazard.

The re-emergence of vanadium as an environmental hazard

Global anthropogenic releases to soil and water for V have been recently re-evaluated and the rate at which V is being deposited into the environment is increasing. As a result, V is again accumulating in the environment. In the atmosphere, V has the highest anthropogenic enrichment factor (AEF) of all trace elements (1) and has the fourth highest AEF in global rivers, behind Cd, Sb and Ni (2). This dominant anthropogenic signal reflects dispersed and pervasive environmental releases in the global V cycle as a result of changing societal demands.

Global V production has approximately doubled in the last 15 years to 80,000 t y⁻¹ in 2017 (3), driven by increased demand for high grade steel. Emerging policy, in the People's Republic of China (policy number: GB/T 1499.2-2018), to increase V content in steel to improve its tensile qualities is expected to increase China's V consumption by 10,000 t y⁻¹. This was reflected in a 100% increase in mined V prices in 2017. There has been a global rise in discharges to the environment of V-rich industrial by-products including steel slags, ash from the expansion of waste incineration (e.g. in the European Union, EU), and bauxite processing residue which now reaches 120 million t y⁻¹ globally (4). Emerging technologies are forecast to enhance global V production and environmental releases. Vanadium redox-flow batteries are being rapidly developed for power storage having the advantage of being able to charge and discharge simultaneously, making them ideal for use in off-grid locations to support renewable energy needs.

Addressing knowledge gaps in vanadium environmental behaviour

Despite the increasing prevalence of V in the environment, we still possess a relatively poor understanding of V geochemistry, relative to other contaminants. Vanadium has three stable oxidation states: V⁺³, V⁺⁴ and V⁺⁵, although it is most commonly found as V⁺⁴ or V⁺⁵, with the latter showing greater solubility under oxic conditions. Vanadium has historically been regarded as a conservative element in surface environments, although there is growing evidence of greater mobility. In freshwater streams affected by the release of red mud in Ajka, Hungary, V exhibited cycling and attenuation behaviour with other ubiquitous elements e.g. aluminum, iron, and molybdenum. These results confirmed that V can disperse and persist in the environment to a greater degree than other contaminants such as arsenic or phosphorus. Vanadium exhibits multiple interactions within surface environments (5) including complexing to organic and inorganic matter in sediments and uptake into flora and fauna, some bacteria being known to scavenge V from refractory compounds. However, the complicated interactions between these processes, and responses to changes in chemical and physical conditions within environmental compartments are poorly understood.

Predicting the fate and behaviour of V in the environment requires that we understand its speciation and phase association. Powerful analytical methods, such as high-resolution transmission electron microscopy and synchrotron-based X-Ray spectroscopy, can provide molecular scale geochemical characterisation. However, these methods require concentrated samples and may not be applicable beyond highly contaminated materials. For the wider environment, methods more suited to lower concentration samples are required, such as ion chromatography-mass spectrometry for aqueous speciation and novel V-specific sequential extractions to understand solid partitioning. The potential to utilise $^{50/51}\text{V}$ isotopic fractionation to trace V through environmental compartments represents an exciting opportunity to assess V behaviour and transport through ecosystems. Data on V in freshwater and sediment monitoring databases (e.g. the US Geological Survey and the Environmental Protection Agency) may be exploited to help describe regional distributions and trends of V in soils, sediments and waters. It is essential that these data are produced to underpin the development and validation of much needed geochemical models to support prediction of environmental risk and behaviour.

A call for preventative measures

We face an increasing likelihood of acute exposure to V that is largely unregulated (6). Some jurisdictions are now remedying this regulatory oversight although much is still to be achieved. In the USA, V is now on the Contaminant Candidate List 4 (CCL4) and is subject to more stringent monitoring in potable waters. Such regulatory attention is encouraging and needs to be adopted more broadly alongside measures to minimise environmental V release. For example, since current global recycling rates for V are estimated by UNEP at <1% (7), there is significant scope for V re-use and/or recycling to meet escalating anthropogenic demand and reduce environmental exposure.

Vanadium hazard and risk assessments must be improved. In the most comprehensive study to date, the W.H.O. concluded that V concentrations in environmental media are substantially lower than toxic concentrations reported in ecotoxicology studies, noting that the paucity of data from specific industrial sites prevented an accurate risk assessment. A review of ecotoxicology data commissioned by the Netherlands' Government has subsequently proposed water quality standards for dissolved V of 1.2 and 3.0 $\mu\text{g L}^{-1}$ for long- and short-term exposure, respectively (8). These standards are similar to the reported background range of concentrations. For example, V concentrations in a large proportion of EU surface waters (range <0.05 $\mu\text{g L}^{-1}$ to 19.5 $\mu\text{g L}^{-1}$; median 0.46 $\mu\text{g L}^{-1}$) (9) exceed or are near to these proposed standards, suggesting that any further increase in V losses to the environment will cause, at least, a major regulatory concern.

The emerging V sources described above, and the legacy of historic emissions represent a growing problem requiring wide scale intervention. The International Aluminium Institute has produced best practice guidance on the management of V-enriched bauxite residues to reduce the likelihood of un-controlled discharges on a global scale. There is a need for other industries to do the same. The global life-cycle of V must be comprehensively mapped and used to identify priority actions through which more sustainable V use can be achieved.

References

1. Schlesinger, W.H., Klein, E.M. and Vengosh, A. Global biogeochemical cycle of vanadium. *Proceedings of the National Academy of Sciences*, **2017** p.201715500.

111 2. Viers, J.; Dupré, B.; Gaillardet, J. Chemical composition of suspended sediments in World
 112 Rivers: New insights from a new database. *Science of the Total Environment*, **2009**, 407: 853-
 113 868.

114 3. United States Geological Survey. Vanadium Statistics and Information. (2018). Available at
 115 <https://minerals.usgs.gov/minerals/pubs/commodity/vanadium/mcs-2018-vanad.pdf> (last
 116 accessed: 05.17.18)

117 4. Power, G., Gräfe, M.; Klauber, C. Bauxite residue issues: I. Current management, disposal
 118 and storage practices. *Hydrometallurgy*, **2011**, 108, 33-45.

119 5. Huang, J.H.; Huang, F.; Evans, L.; Glasauer, S. Vanadium: Global (bio) geochemistry.
 120 *Chemical Geology*, **2015**, 417, 68-89.

121 6. Rattner, B.A.; McKernan, M.A.; Eisenreich, K.M.; Link, W.A.; Olsen, G.H.; Hoffman, D.J.;
 122 Knowles, K.A.; McGowan, P.C. Toxicity and hazard of vanadium to mallard ducks (*Anas*
 123 platyrhynchos) and Canada geese (*Branta canadensis*). *Journal of Toxicology and*
 124 *Environmental Health*, **2006**, 69, 331-351.

125 7. Graedel, T.E.; Allwood, J.; Birat, J.P. ; Buchert, M. ; Hagelüken, C. ; Reck, B.K. ; Sibley,
 126 S.F. ; Sonnemann, G. Recycling rates of metals: A status report, *United Nations Environment*
 127 *Programme*, **2011**

128 8. Smit, C.E. Environmental Risk Limits for Vanadium in Water: A Proposal for Water Quality
 129 Standards in Accordance with the Water Framework Directive. RIVM Letter Report
 130 601714021/2012. **2012** National Institute for Public Health and the Environment, Bilthoven,
 131 73.

132 9. Salminen, R. ; Batista, M.J. ; Bidovec, M. ; Demetriades, A. ; De Vivo, B. ; De Vos, W. ;
 133 Duris, M. ; Gilucis, A. ; Gregorauskiene, V. ; Halamić, J. ; Heitzmann, P. 2005. Geochemical
 134 atlas of Europe, part 1, background information, methodology and maps. Geological survey of
 135 Finland.